arXiv:1210.3737v1 [cond-mat.stat-mech] 13 Oct 2012

A non-classical van der Waals loop: Collective variables method

R V Romanik and M P Kozlovskii

Institute for Condensed Matter Physics of the National Academy of Sciences of Ukraine, 1 Svientsitskii Str., 79011 Lviv, Ukraine

E-mail: `romanik@icmp.lviv.ua`

Abstract. The equation of state is investigated for an Ising-like model in the framework of collective variables method. The peculiar feature of the theory is that a non-classical van der Waals loop is obtained. The results are compared with ones of a trigonometric parametric model, in terms of normalized magnetization, $\tilde{M}$, and field, $\tilde{H}$.

PACS numbers: 05.50.+q, 05.70.Ce, 64.60.F-, 75.10.Hk

1. Introduction

Recently, in [1] it was shown that considering the system of Ising spins on a simple cubic lattice by the collective variables (CV) method [2, 3], a non-classical van der Waals (vdW) loop can be obtained. In the present paper, the shape of the loop is investigated. A comparison is made with similar results of [4], where the van der Waals loop was obtained for a system with a scalar order parameter using a trigonometric parametric model for scaling behaviour near criticality. The discussion on whether such a loop exists naturally and has or not any physical manifestation can be found in [5]. Our purpose in this paper is to compare results from CV theory with others.

We consider a system of N Ising spins on a simple cubic lattice of spacing c. The Hamiltonian of such a system is well known

$$H = -\frac{1}{2}\sum_{\boldsymbol{i},\boldsymbol{j}} \Phi(r_{\boldsymbol{i},\boldsymbol{j}})\sigma_{\boldsymbol{i}}\sigma_{\boldsymbol{j}} - \mathcal{H}\sum_{\boldsymbol{i}} \sigma_{\boldsymbol{i}}. \qquad (1)$$

Here the spin variables σ_i take on ±1, $\mathcal{H}$ is the external field, and $\Phi(r_{\boldsymbol{i},\boldsymbol{j}})$ is a short-range interaction potential between spins located at the i-th and j-th sites of separation r_{ij}. The interaction potential can be chosen in the form of exponentially decreasing function, $\Phi(r_{ij}) = const \cdot \exp(-r_{ij}/b)$, with b being an effective range.

The partition function

$$Z = \sum_{\{\sigma\}} \mathrm{e}^{-\beta H}, \qquad (2)$$

where $\beta = (k_{\mathrm{B}}T)^{-1}$ is the inverse temperature, can be written in terms of collective variables $\rho_{\mathbf{k}}$ [2, 3]. In "ρ^4-model" approximation, the explicit form for such a representation is as follows

$$Z = Z_0 \int (\mathrm{d}\rho)^{N_0} \exp\Big[a_1\sqrt{N_0}\rho_0 - \frac{1}{2}\sum_{\mathbf{k}\in\mathcal{B}_\mathbf{0}} d(k)\rho_{\mathbf{k}}\rho_{-\mathbf{k}} - - \frac{a_4}{4!}N_0^{-1}\sum_{\mathbf{k}_i\in\mathcal{B}_0} \rho_{\mathbf{k}_1}\dots\rho_{\mathbf{k}_4}\delta_{\mathbf{k}_1+\ldots+\mathbf{k}_4}\Big]. \qquad (3)$$

Here the quantity $d(k)$ contains the Fourier transform of the interaction potential

$$d(k) = a_2 + \beta\Phi(0)\bar{\Phi} - \beta\Phi(k). \qquad (4)$$

For $\tilde{\Phi}(k)$ we use the so-called parabolic approximation

$$\tilde{\Phi}(k) = \begin{cases} \Phi(0)(1-2b^2k^2) & \text{for } k \le B_0 = Bs_0^{-1} \\ \Phi_0 = \Phi(0)\bar{\Phi} & \text{for } B_0 < k \le B. \end{cases} \qquad (5)$$

Strictly speaking, the wave vector $\mathbf{k}$ takes on values from the first Brillouin zone

$$\mathcal{B} = \{\mathbf{k} = (k_x, k_y, k_z) | k_i = -\frac{\pi}{c} + \frac{2\pi}{c}\frac{n_i}{N_i}; n_i = 1, 2, \dots, N_i; i = x, y, z\}. \quad (6)$$

In the following, however, we will keep to the spherical approximation for the Brillouine zone so that $B = \pi/c$ is the boundary of this zone, and $B_0 = \pi/(cs_0) = \pi/c_0$ is the boundary of $\mathcal{B}_0$. The discussion on the choice for s_0 and $\bar{\Phi}$ can be found in [6]. In general, s_0 should depend on the ratio of the effective interaction range b to the lattice constant c. However, in the present calculation, we fix $s_0 = 2$ and $b/c = 0.3$. This gives $\bar{\Phi} = 0.27492$. The quantities a_i from (3) are expressed as

$$a_1 = s_0^{d/2} h, \qquad a_2 = 1 - s_0^{-d}, \qquad a_4 = 2 s_0^{-d}, \tag{7}$$

where $d = 3$ is the space dimension, and $h = \beta\mathcal{H}$ is dimensionless field. In (3) the collective variables $\rho_{\mathbf{k}}$ with $B_0 < k \leq B$ have already been integrated out so that $N_0 = N s_0^{-3}$ is a number of variables remaining to be integrated.

Calculation of the partition function (3) is performed according to Yukhnovskii's method [2]. It is based on the idea of step-by-step integration of the partition function, over the subsets of collective variables, first with $B_1 < k \leq B_0$, then with $B_2 < k \leq B_1$, and so on, while averaging the Fourier image of the interaction potential on each step. Here $B_n = B_{n-1}/s = B_0/s^n$, where s is the renormalization group (RG) parameter. This is equivalent to the Kadanoff scheme of constructing spin blocks [7, 8]. Every time when integrating over a subset of CV, a factor – denote it Q_n – appears in the partition function. On performing step-by-step integration of the partition function over n_{p} subsets, one arrives at

$$Z = Z_0 [Q(d)]^{N_0} \Big(\prod_{n=1}^{n_{\mathrm{p}}} Q_n \Big) Z_{\mathrm{LGR}}. \tag{8}$$

The partial partition functions Q_n of the n-th level are characterized by a set of coefficients $a_1^{(n)}$, $d_n(0)$, $a_4^{(n)}$, for which the recurrence relations (RR) hold [9]. The RR have a fixed point as a partial solution. Because of that, the quantity n_{p} is chosen from the requirement that for $n \leq n_{\mathrm{p}}$ the RR can be linearized near the fixed point. In this case, the system possesses the RG symmetry and is said to be in the critical regime of the order parameter fluctuations. The quantity n_{p} is called the exit point from the critical regime.

The analytic expressions for the exit point were found for the limiting cases, $h = 0$ and $\tau = 0$:

$$\begin{aligned} n_{\mathrm{p}} &= -\frac{\ln|\tilde{h}|}{\ln E_1} - 1, \qquad \text{at} \quad \tau = 0, \\ &= -\frac{\ln|\tilde{\tau}|}{\ln E_2} - 1, \qquad \text{at} \quad h = 0, \end{aligned} \tag{9}$$

where $\tau = (T - T_{\mathrm{c}})/T_{\mathrm{c}}$ is the reduced temperature.

Here $E_1 = 24.551$ and $E_2 = 8.306$ are the eigenvalues of the matrix of the RG transformation linearized near the fixed point of RR,

$$\tilde{\tau} = \frac{c_{1\mathrm{k}}}{f_0} \tau, \qquad \tilde{h} = \frac{s_0^{d/2}}{h_0} h, \tag{10}$$

where $f_0 = 0.5$ defines the fixed-point coordinates, $c_{1\mathrm{k}} = 1.155$, $h_0 = 0.760$. In general case, the expression for n_{p} cannot be obtained analytically. For example, in [10] this quantity was computed numerically as a solution to a certain equation. In any case, n_{p}

should satisfy (9) in the mentioned limiting cases. Based on this requirement, in [11] the expression was constructed for the exit point in the form

$$n_{\mathrm{p}}^{(\pm)} = -\frac{\ln(\tilde{h}^2 + h_{\mathrm{c}}^{(\pm)^2})}{2\ln E_1} - 1, \tag{11}$$

where some temperature fields, $h_{\mathrm{c}}^{(+)} = |\tau|^{\beta\delta}$ and $h_{\mathrm{c}}^{(-)} = |\tau_1|^{\beta\delta}$, are introduced, β and δ are the critical exponents of magnetization. In our approach, $\beta = 0.3024$ and $\delta = d+2 = 5$. The signs "+" and "-" are related to $T > T_{\mathrm{c}}$ and $T < T_{\mathrm{c}}$, respectively. In the following, we will mainly omit superscript ±. Finally, $\tau_1 = -E_2^{n_0}\tau$, where n_0 denotes the difference between $n_{\mathrm{p}}^{(+)}$ and $n_{\mathrm{p}}^{(-)}$. In [12] $n_0 = 0.5$ is chosen to recover the universal ratio of critical amplitudes for the correlation length, $\xi^+/\xi^- = 1.896(10)$ [13].

In (8), the quantity Z_{LGR} is still remaining undefined. Here, we shall not delve into considering this quantity. A detailed explanation of how to compute it can be found in [12, 14]. We just recall that it is expressed in the form

$$Z_{\mathrm{LGR}} = Z_{\mathrm{G}} Z_{\mathrm{TR}} \cdot \mathrm{e}^{NE_0(\sigma)}, \tag{12}$$

where the contribution $e^{NE_0(\sigma)}$ is the most important because it is due to the collective variable ρ_0. In the present research, main attention is paid on this part of the partition function. The quantity $E_0(\sigma)$ has the form

$$E_0 = e_0 h(\tilde{h}^2 + h_{\mathrm{c}}^{(\pm)^2})^{\frac{1}{2(d+2)}} - e_2(\tilde{h}^2 + h_{\mathrm{c}}^{(\pm)^2})^{\frac{d}{d+2}}. \tag{13}$$

The following notation is used

$$\begin{aligned} e_0 &= \sigma_0 s^{-1/2}, \\ e_2 &= \frac{1}{2}\sigma_0^2 s^{-3}(r_{n_{\mathrm{p}}+2} + \frac{1}{12}s_0^3\sigma_0^2 u_{n_{\mathrm{p}}+2}), \end{aligned} \tag{14}$$

where

$$\begin{aligned} r_{n_{\mathrm{p}}+2} &= \beta\Phi(0) f_0(-1 \pm E_2 H_{\mathrm{c}}), \\ u_{n_{\mathrm{p}}+2} &= (\beta\Phi(0))^2 \varphi_0(1 \pm \Phi E_2 H_{\mathrm{c}}), \end{aligned} \tag{15}$$

with $H_{\mathrm{c}} = \tilde{\tau}(\tilde{h}^2 + h_{\mathrm{c}}^{(\pm)^2})^{-1/2p_0}$. The quantity σ_0 is the solution of the following cubic equation

$$\sigma_0^3 + p\sigma_0 + q = 0, \tag{16}$$

with the coefficients

$$\begin{aligned} p &= 6s_0^{-3}\frac{r_{n_{\mathrm{p}}+2}}{u_{n_{\mathrm{p}}+2}}, \\ q &= -6s_0^{-9/2}s^{5/2}\frac{h_0}{u_{n_{\mathrm{p}}+2}}\frac{\tilde{h}}{(\tilde{h}^2 + h_{\mathrm{c}}^{(\pm)^2})^{1/2}}. \end{aligned} \tag{17}$$

2. Equation of state

Based on (8) and (12), the Gibbs free energy can be expressed as a sum of three terms

$$F(\tau,h) = -k_{\mathrm{B}}T\ln Z = F_{\mathrm{a}} + F_{\mathrm{s}}^{(\pm)} + F_0^{(\pm)}. \tag{18}$$

Here, the term F_{a} is the analytical part of the free energy and does not affect the critical behaviour. The term $F_{\mathrm{s}}^{(\pm)}$ is expressed as

$$F_{\mathrm{s}}^{(\pm)}(\tau,h) = -k_{\mathrm{B}}TN\gamma_{\mathrm{s}}^{(\pm)}(\tilde{h}^2 + h_{\mathrm{c}}^{(\pm)^2})^{\frac{d}{d+2}}, \tag{19}$$

where $\gamma_{\mathrm{s}}^{\pm}$ includes contributions from the critical regime and from the limiting Gaussian regime (inverse Gaussian regime in the case of $T < T_{\mathrm{c}}$). The explicit expression for it can be found in (5.6) of [14]. Finally, the quantity $F_0^{(\pm)}$ from (18) is

$$F_0^{(\pm)}(\tau,h) = -k_{\mathrm{B}}TNE_0(\sigma). \tag{20}$$

This contribution to the Gibbs free energy is due to the collective variable ρ_0, which, as is known from the theory of collective variables [2], is related to the order parameter. Therefore, $F_0^{(\pm)}$ is the free energy of ordering. The order parameter of the considered system – the magnetization – is then calculated by means of the thermodynamic formula

$$M_0 = -\frac{1}{N}\frac{\partial F_0^{\pm}}{\partial\mathcal{H}} = \frac{\partial E_0(\sigma)}{\partial h} \tag{21}$$

that leads to the expression

$$M_0 = (\tilde{h}^2 + h_{\mathrm{c}}^2)^{\frac{1}{2(d+2)}}\Bigg[e_0\Bigg(1 + \frac{1}{5}\frac{\tilde{h}^2}{\tilde{h}^2 + h_{\mathrm{c}}^{(\pm)^2}}\Bigg) - \frac{6}{5}e_2\frac{s_0^{3/2}}{h_0}\frac{\tilde{h}}{(\tilde{h}^2 + h_{\mathrm{c}}^{(\pm)^2})^{1/2}} - \\ - \left(\frac{\mathrm{d}e_2}{\mathrm{d}h}\right)_{\sigma}(\tilde{h}^2 + h_{\mathrm{c}}^{(\pm)^2})^{1/2}\Bigg]. \tag{22}$$

The magnetization (22) can be rewritten in more compact form

$$M_0 = \sigma_{00}(\tilde{h}^2 + h_{\mathrm{c}}^{(\pm)^2})^{\frac{1}{2\delta}}, \tag{23}$$

where $\delta = 5$ is the critical exponent describing the field dependence of magnetization and

$$\sigma_{00} = \Bigg[e_0\Bigg(1 + \frac{1}{5}\frac{\tilde{h}^2}{\tilde{h}^2 + h_{\mathrm{c}}^{(\pm)^2}}\Bigg) - \frac{6}{5}e_2\frac{s_0^{3/2}}{h_0}\frac{\tilde{h}}{(\tilde{h}^2 + h_{\mathrm{c}}^{(\pm)^2})^{1/2}} - \\ - \left(\frac{\mathrm{d}e_2}{\mathrm{d}h}\right)_{\sigma}(\tilde{h}^2 + h_{\mathrm{c}}^{(\pm)^2})^{1/2}\Bigg]. \tag{24}$$

In figure 1 the magnetization (22) is presented as a function of the external field. As is seen from the picture, the proposed approach gives a van der Waals loop, however with non-classical exponents. In order to compare our results with ones obtained by different methods, we appeal to [4], where a non-classical van der Waals loop was obtained with the help of a trigonometric parametric model for scaling behaviour near criticality. For this reason, in the next section the main features of the conventional parametric presentations of the equation of state are presented as well as the novelty suggested in [4].



Figure 1. Magnetization (22) as a function of field below the critical temperature, $\tau = -0.001$.

3. Parametric representation of the equation of state

The way of representation of the critical behaviour with the aid of new parameters, r and θ, was firstly suggested in [15]. The meaning of the parameters is that the "radial" variable r defines the distance from the critical point on the "temperature-field" plane, and the "angular" variable θ defines the path along a contour of constant r. Today, the more traditional form of the parametric representation is as follows

$$M = r^{\beta} m(\theta), \qquad h = r^{\beta\delta} l(\theta), \qquad \tau = r k(\theta), \tag{25}$$

where, in terms of magnet, M is the magnetization, h is the magnetic field, and τ is the reduced temperature.

The function $k(\theta)$ is usually chosen in the form $k(\theta) = 1 - b^2\theta^2$, which means that $\theta_{\mathrm{c}} = \pm 1/b$ corresponds to the critical isotherm, for $h > 0$ and $h < 0$, respectively. The function $l(\theta)$ is chosen in the form of a polynomial with only odd powers of θ. Zeros of $l(\theta)$ correspond to the absence of the field, $h = 0$. Particularly, $\theta = 0$ specifies the critical isochore, $h = 0$ and $\tau = r > 0$. Next zeros of $l(\theta)$, $\theta = \pm\theta_1$, correspond to two sides of the coexistence curve. $l(\theta)$ is usually written in the form $l(\theta) = l_0 l_1(\theta)$ so that $\mathrm{d}l(\theta)/\mathrm{d}\theta|_{\theta=0} \equiv l'(0) = l_0$, and l_0 is a normalization constant. And, finally, $m(\theta) = m_0(\theta + c\theta^3)$. If $c = 0$, the transformation (25) is referred to as the "linear" model, and in the case of $c \neq 0$ it is called the "cubic" model.

As can be easily seen, such a representation fails in the two phase region. For this reason, in [5, 4] were suggested new, the so-called trigonometric parametric models that would naturally extend to the two phase region and recover a van der Waals loop. Here we present the main features of the extended sine model [4] – the more preferable of the trigonometric models.

First of all, in this model a transformation of type (25) touches the temperature, magnetization and the thermodynamic potential $\mathcal{A}(M,T)$:

$$\tau = rk(\theta), \qquad M = r^{\beta}m(\theta), \qquad \mathcal{A} = r^{2-\alpha}n(\theta), \tag{26}$$

where the angular functions are chosen in the form

$$\begin{aligned} k(\theta) &= \ 1 - 2b^2[1-\cos(q\theta)]/q^2, \\ m(\theta) &= m_0 \sin(q\theta)/q, \\ n(\theta) &= \ l_0 m_0 [1 + \sum_{i=1}^{j} c_i k^i(\theta)]. \end{aligned} \tag{27}$$

Here the coefficients b, q, and c_i are regarded as adjustable parameters. In such circumstances, for the field angular function $l(\theta)$ one has

$$l(\theta) = l_0 \frac{k(\theta)n_1'(\theta) - (2-\alpha)k'(\theta)n_1(\theta)}{k(\theta)m_1'(\theta) - \beta k'(\theta)m_1(\theta)}, \tag{28}$$

where we have introduced the following notation

$$\begin{aligned} n_1(\theta) &= \ \frac{n(\theta)}{l_0 m_0} = 1 + \sum_{i=1}^{j} c_i k^i(\theta), \\ m_1(\theta) &= \frac{m(\theta)}{m_0} = \sin(q\theta)/q. \end{aligned} \tag{29}$$

The van der Waals loop is obtained in terms of normalized versions of M and h, namely

$$\tilde{M} = M/M_0(T), \qquad \tilde{H} = h/[M_0(T)/\chi_0^+(T)], \tag{30}$$

where $M_0(T)$ is the "spontaneous" value of the order parameter, and $\chi_0^+(T)$ is the zero-field susceptibility above T_c, $\chi_0^+(T) \approx C^+\tau^{-\gamma}$.

In figure 2, the dashed line is the van der Waals loop for the following values of the adjustable parameters (see formula (6.2) in [4]):

$$\begin{aligned} &b = 3.81857, &&q = 3.03134, &&c_1 = 1.71886 \\ &c_2 = -5.97266, &&c_3 = -0.207431, &&c_4 = 0.0882037. \end{aligned} \tag{31}$$

Note, that for such a choice $l'(0) \neq l_0$, in contrast to either the "linear" or "cubic" model.

In the same figure 2, the solid line is our results based on (22). Regarding the approximations, "ρ^4-model", and zero value of the small critical exponent $\eta = 0$, we see good qualitative agreement in results, especially in the regions of $\tilde{h} \approx 0$ and of $\tilde{h} \gg h_{\mathrm{c}}$, $\tilde{H} > 2$. Some qualitative discrepancy is observed for the intermediate values of $\tilde{H}$, i.e., for $\tilde{h} \approx h_{\mathrm{c}}$. In this domain, with increasing $\tilde{H}$ our curve initially withdraws from the dashed curve, further approaches and intersects it and then moves along with the dashed curve, below it all the time. Such discrepancy is connected with the choice of functional form of n_{p} (11). Although it gives correct values in the limiting cases, see (9), however it seems to fail in the intermediate region, $\tilde{h} \approx h_{\mathrm{c}}$. Therefore, to improve our results we



Figure 2. Scaled van der Waals loop and magnetization above the critical temperature. The dashed curves represent results obtained within the extended sine model. The full curves represent the results of the collective variables method, equation (22).

need a somewhat different functional form in this value range. Another way to do so is to compute n_{p} numerically, but we do not lose hope to solve the problem analytically and will attempt to find more appropriate expression for the exit point in a future work.

With regard to the loop, we observe the loop is wider in CV theory. The spinodal value of the magnetization in CV, $\tilde{M} = 0.55$ is less then the corresponding value $\tilde{M} = 0.697$ from [5], which means that the CV theory, at least in "ρ^4-model" approximation, gives wider metastable domain.

4. Conclusions

In conclusion, this work is the first attempt to investigate the van der Waals loop using the collective variables method, which is essentially microscopic approach. This investigation is important because there is lack of non-classical theories that give the van der Waals loop (see, e.g., [16], Sec. 3.4.4). Some quantitative disagreement of our result in comparison to ones obtained in [4] can be associated with the model approximation, since present calculations are carried out using the simplest non-Gaussian approximation, i.e., ρ^4-model. The obtained result can be improved both in a formal way, by appropriately choosing the functional form of the exit point n_{p}, and in a conceptual way, to which can be attributed (a) investigating the system in a higher approximation, i.e., ρ^n-model with $n > 4$ (for $n = 6$ see [17]), (b) including corrections to scaling [18], and/or (c) accounting for averaging the interaction potential [19], the latter resulting in $\eta \neq 0$. All of these will be the scope of a forthcoming paper.

References

[1] Kozlovskii M P and Romanik R V 2011 *Condens. Matter Phys.* **14** 43002

[2] Yukhnovskii I R 1987 *Phase Transitions of the Second Order. Collective Variables Method* (Singapore: World Scientific) p 224
[3] Yuchnovskii I R, Kozlovskii M P and Pylyuk I V 2001 *Microscopic Theory of Phase Transitions in the Three Dimensional Systems* (Lviv: Eurosvit) p 592 (in Ukrainian)
[4] Fisher M E, Zinn S-Y and Upton P J 1999 *Phys. Rev.* B **59** 14533
[5] Fisher M E and Zinn S-Y 1998 *J. Phys. A: Math. Gen.* **31** L629
[6] Kozlovskii M P and Romanik R V 2010 *Condens. Matter Phys.* **13** 43004
[7] Kadanoff L P 1966 *Physics (N.Y.)* **2** 263
[8] Kadanoff L P 1967 *Rev. Mod. Phys.* **39** 395
[9] Kozlovskii M P 2005 *Condens. Matter Phys.* **8** 473
[10] Kozlovskii M P, Pylyuk I V and Prytula O O 2006 *Nucl. Phys.* B **753** 242
[11] Kozlovskii M P 2007 *Phase Transitions* **80** 3
[12] Kozlovskii M P 2009 *Ukr. Fiz. Zh./Reviews (Ukr. ed.)* **5** 61 (in Ukrainian)
[13] Campostrini M, Pelissetto A, Rossi P and Vicari E 2002 *Phys. Rev.* E **65** 066127
[14] Kozlovskii M P 2009 *Condens. Matter Phys.* **12** 151
[15] Schofield P 1969 *Phys. Rev. Lett.* **22** 606.
[16] Pelissetto A and Vicari E 2002 *Phys. Rept.* **368** 549
[17] Kozlovskii M P and Pylyuk I V 2011 *ISRN Condens. Matter Phys.* 260750
[18] Yukhnovskii I R, Kozlovskii M P and Pylyuk I V 2002 *Phys. Rev.* B **66** 134410
[19] Yukhnovskii I R, Kozlovskii M P and Pylyuk I V 2012 *Ukr. J. Phys.* **57** 80